

# The Relationship Between Childhood Abuse and Adult Attachment Styles: The Mediator Role of Sensory Over-Responsivity

Alexandra Silva<sup>a,b</sup>, Simão Ferreira<sup>a</sup>, Élia Silva Pinto<sup>b</sup>, Susana Aguiar Rocha<sup>c</sup>, and Nuno Barbosa-Rocha<sup>a</sup>

<sup>a</sup>Center for Translational Health and Medical Biotechnology Research, School of Health, Polytechnic of Porto, Porto, Portugal; <sup>b</sup>Occupational Therapy Department, Alcoitão School of Health, Alcabideche, Portugal; <sup>c</sup>Porto Accounting and Business School, Polytechnic of Porto, Porto, Portugal

## ABSTRACT

Child abuse portrays health and well-being issues that can last for several years, including altering interpersonal behavior. Evidence has shown that these early negative experiences may cause changes in sensory modulation. This study aimed to understand if sensory over-responsivity (SOR) plays an important role in mediating the association between childhood abuse (emotional abuse, emotional neglect, sexual abuse, physical abuse, physical neglect) and attachment-related anxiety or avoidance in adult romantic relationships. An online survey was conducted to examine these associations in a Portuguese community sample ( $N = 500$ ) aged 18–62 years. The presented mediation models shown a higher mediation percentage for attachment-related anxiety rather than avoidance. All variables predicted the mediator, except sexual abuse which did not predict the overall model when the mediator was included. Our results have implications for individual treatment approaches regarding child abuse victims and can certainly influence intervention strategies.

## ARTICLE HISTORY

Received 17 February 2022

Revised 2 February 2023

Accepted 4 February 2023

## KEYWORDS

Attachment; sensory over-responsivity; traumatic childhood experiences

Child abuse is a persistent problem in society and affects many children around the world. A review concluded that, worldwide, the overall estimated prevalence of childhood maltreatment was 127/1000 for sexual abuse, 226/1000 for physical abuse, 363/1000 for emotional abuse, 163/1000 for physical neglect, and 184/1000 for emotional neglect (Stoltenborgh et al., 2015). According to the Portuguese Association of Victim Support (Associação

**CONTACT** Nuno Barbosa-Rocha  [nrocha@ess.ipp.pt](mailto:nrocha@ess.ipp.pt)  Center for Translational Health and Medical Biotechnology Research, School of Health, Polytechnic of Porto, Rua Dr. António Bernardino de Almeida, Porto 400 4200 - 072, Portugal

Alexandra Silva and Simão Ferreira share first co-authorship as they equally contributed to the paper.

Author note

Silva A., Center for Translational Health and Medical Biotechnology Research, School of Health, Polytechnic of Porto, Porto, Portugal and Alcoitão School of Health, Alcabideche, Portugal; Ferreira S., Center for Translational Health and Medical Biotechnology Research, School of Health, Polytechnic of Porto, Porto, Portugal; Pinto E., Alcoitão School of Health, Alcabideche, Portugal; Rocha S., ISCAP, Polytechnic of Porto, Porto, Portugal; Barbosa-Rocha N., Center for Translational Health and Medical Biotechnology Research, School of Health, Polytechnic of Porto, Porto, Portugal.

© 2023 The Author(s). Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

Portuguesa de Apoio à Vítima, APAV), there were 1841 abused children and adolescents in Portugal in 2020, of which 59.7% are female (APAV, 2021). These numbers are probably an underrepresentation of the reality. Determining the exact incidence of child abuse and neglect is challenging, mostly because cases are frequently underreported or unnoticed (Gubbels et al., 2021).

Child abuse or maltreatment constitutes all forms of physical and/or emotional ill-treatment, sexual abuse, neglect or negligent treatment, and exploitation, resulting in actual or potential harm to the child's health, survival, development or dignity in the context of a relationship of responsibility, trust or power (WHO, 2022). Consequences include health problems, social and behavioral problems (mainly, attachment problems) impaired cognitive and academic performance and economic problems.

In short, physical abuse is any action that causes, or has the potential to cause, physical harm to a child. Emotional abuse includes not providing an appropriate and supportive environment and having negative attitudes toward the child. Child sexual abuse involves a child being involved in sexual activity that they are not prepared for or do not understand. Neglect is when a child's development is not supported in areas like health, education, and emotional development, despite resources being available. Commercial or other exploitation is when a child is used for someone else's benefit, such as in work (WHO, 1999).

Neglect, one of the forms of child maltreatment, can be classified into two types: physical neglect and emotional neglect. Physical neglect occurs when a caregiver fails to provide a child with basic physical necessities such as food, clothing, shelter, personal hygiene, and medical care. Emotional neglect, on the other hand, refers to a caregiver's failure to meet a child's developmental or emotional needs, such as inadequate nurturance or affection (Cohen et al., 2017).

These early traumatic experiences can negatively impact health outcomes, including long-term mental health problems such as post-traumatic stress disorder, depression, anxiety disorders, alcohol problems, antisocial behavior, aggressive behavior, anger, suicidal behavior, and persistence of medically unexplained physical symptoms (Christ et al., 2019). Moreover, these negative experiences may have a lasting impact through adulthood in the way those that suffered childhood abuse relate to other people (Rueness et al., 2020).

According to attachment theory, infancy is the time to develop cognitive models of relationships with others based on interactions with early caregivers, usually a parent or parents. Children whose needs are met, develop personal models of being competent, effective, and lovable, promoting self-esteem and coherent ego development; models of other people being predictable and trustworthy; and models of relationships as potentially rewarding and worthwhile. According to attachment theory, these children also experience a sense

of security and readiness to explore the environment, while maintaining parental figures as a safety-net to whom they can return if necessary (Bowlby, 1982). Childhood traumatic experiences regarding caregivers, are expected to evolve into insecure emotional attachment styles that are reflected in relationships with others during adolescence and even in adulthood (Unger & De Luca, 2014).

Maltreated children, especially those who have been physically abused, have been reported to have a less positive self-concept than others (Arata et al., 2005). Children who are neglected and maltreated may build relationship models characterized by a lack of responsiveness. They may withdraw from social relationships, develop relationships characterized by mistrust, leading to aggressive or defensive behaviors, that may engage in poorly controlled behavior leading to being excluded from relationships (Bacon & Richardson, 2001). Similarly, child abuse experiences have a detrimental impact on future adult romantic relationships and attachment styles (Yoon, 2020). Although the risk for negative interpersonal outcomes is high, counterintuitively it seems that the experience of childhood abuse may bring an opportunity for personal growth, prosociality, and increased empathy, that may imply different mechanisms and mediators (Greenberg et al., 2018).

Some studies found that those who have experienced physical, sexual, and emotional abuse have been related to negative functioning in adult romantic relationships (McCarthy & Taylor, 1999). Other studies suggest that participants who experienced abuse showed more insecure attachment styles than participants who had not experienced abuse (Unger & De Luca, 2014).

An intimate relationship with a partner can be classified as a secure, avoidant (withdrawn), and anxious (worried) attachment. Thus, a secure attachment in intimate relationships is characterized by trust, friendship, satisfaction, reciprocity, self-disclosure, commitment, and guidance in joint problem solving. An individual with an avoidant attachment does not appear to be invested in intimate relationships and avoids the partner in various situations; and with said anxious attachment, they may find that it is difficult to be comforted by the partner because of the uncertainty associated with insecurities (insecurity about being loved and protected, insecurity about deserving love) (Bolger et al., 1998).

In this study, we hypothesize that sensory over-responsivity (SOR) may play an important role in mediating the association between childhood abuse and attachment-related anxiety and avoidance in adult romantic relationships.


## The role of sensory over-responsivity

Childhood abuse experiences can lead to future sensory challenges (Joseph et al., 2021; McGreevy & Boland, 2020). This could be due to a disorder of sensory modulation, which is a disorder of the regulatory component of sensory processing that disrupts the ability to regulate and organize in a graded manner the magnitude, intensity, and type of response to sensory input (Dunn, 2007). When sensory input is modulated into an adaptive behavior, there should be an appropriate balance between habituation and sensitization. Habituation occurs when the central nervous system recognizes the stimuli as familiar and no longer needs to respond to them. Conversely, when the central nervous system recognizes the stimuli as harmful or important, sensitization occurs, and responding increases (Bar-Shalita & Cermak, 2020).

One of these altered responses to stimulus is referred to as SOR (Miller et al., 2012). SOR is characterized by a tendency to perceive benign sensations as threatening, distracting, or painful. SOR is most commonly characterized by intolerance to sound (auditory stimuli) and/or intolerance to touch (tactile stimuli). In a small number of individuals, negative responses occur in other sensory domains such as taste and smell (Bundy & Lane, 2019). The tendency is to react in a fight/flight manner, manifested as exaggerated avoidant and defensive behaviors that are out of sync with the environmental demands. In childhood, these extreme reactions can impact on the child's ability to perform a range of essential developmental tasks and adaptive functions, including social problem solving and overt signs of empathy (Witte et al., 2020). They can also negatively impact early relationships and positive social participation during school years (McGlone et al., 2014), which can persist into adulthood. The long-term negative effects on social relationships may include the preference for greater interpersonal distance and negative reactions to social touch (Maier, Gieling, et al., 2020).

There is some evidence that abused children experience being overresponsive to tactile sensation and auditory filtering (Joseph et al., 2021). In a neurophysiological level, individuals who have experienced early adversity or trauma show altered neural response to stimuli and hyperactive brain activity in regions that detect information, including the amygdala, the insula, the ventrolateral prefrontal cortex, and the dorsal anterior cingulate cortex, which is consistent with the experience of SOR (Maier, Heinen-Ludwig, et al., 2020). Increased bottom-up detection of stimuli may compete with the ability to recruit higher-order brain systems and may interfere with positive social relationships (Jedd et al., 2015).

Since dysregulation of trauma can alter neurochemistry of the central nervous system and integration of the brain, children who have difficulties processing sensory information often show either under reactions or



**Figure 1.** Conceptual diagram of the total mediation analysis.

overreactions, which is considered hypersensitivity, to the sensations of touch, movement, sight, sound, and smell (Whiting, 2018).

### Overview and hypotheses

The purpose of the current study was to examine if SOR is a mediator between the five types of child abuse (emotional abuse, emotional neglect, sexual abuse, physical abuse, physical neglect) and adult attachment (avoidance and anxiety). The hypothesized moderated mediation model (see Figure 1) was tested in Hayes model 4 (Hayes, 2013), using a bootstrapping approach to assess direct and indirect effects of child abuse mediated by over-responsivity, with bias-corrected 95% confidence intervals ( $n = 5000$ ). Moderated mediation analyses test the effect of a predictor (child abuse) on an outcome variable (adult attachment, anxiety, and avoidance) via the potential mediator (SOR). Significant effects are supported by the absence of zero within the confidence intervals.

### Participants

The sample consisted of 500 individuals aged 18–62 years ( $M = 28.83$ ,  $SD = 7.82$ ). The sample is predominantly female (70.6%), with a higher percentage

**Table 1.** Sample Characteristics/Demographics (*n* = 500).

	Mean ± SD	Min - Max
Age	28.83 ± 7.82	18–62
Gender	<b>Groups</b>	<b>n (%)</b>
	<i>Male</i>	147 (29.4%)
	<i>Female</i>	353 (70.6%)
Marital Status	<i>Single</i>	382 (76.4%)
	<i>Married/Living with a partner</i>	104 (20.8%)
	<i>Widower</i>	1 (0.2%)
	<i>Separated/Divorced</i>	13 (2.6%)
Education	<i>Basic Education</i>	13 (2.6%)
	<i>High School</i>	107 (21.4%)
	<i>Bachelor's Degree</i>	273 (54.6%)
	<i>Master's Degree</i>	104 (20.8%)
	<i>Doctoral</i>	3 (0.6%)
Financial situation	<i>Much less than enough money</i>	71 (14.2%)
	<i>Enough money</i>	321 (64.2%)
	<i>More than enough money</i>	91 (18.2%)
	<i>Much more than enough money</i>	1 (0.2%)

of single individuals (76.4%). Additional demographic information can be found in [Table 1](#).

## Measures

### Sensory processing

The Sensory Processing Scale (now called the Sensory Processing Three Dimensions Scale (SP3D)) consists of two parts: an inventory completed by parents, caregivers, or self, and a performance measure or assessment completed by an examiner (Mulligan et al., 2019). In this study, only the inventory was administered by self-completion. The Sensory Processing Scale Inventory (SPSI) reflects sensory reactivity, including SOR, sensory under-responsivity (SUR), and sensory craving (SC) in several sensory domains. The SOR subscale consists of 80 items, the SUR consists of 30 items, and the SC consists of 38 (Schoen et al., 2017). Total scores are then computed for each subtype, with higher scores reflecting a greater number of atypical sensory symptomatology. Internal consistency was strong for each subscale. Alpha coefficient for the total SOR scale was .89, .88 for the total SUR scale, and .93 for the total SC scale.

### Childhood maltreatment

The Childhood Trauma Questionnaire (CTQ-SF) is a 28-item self-report instrument for abusive situations occurring up to age 15. It is a reduced version of the original questionnaire consisting of 70 items. It is classified on a 5-point Likert scale corresponding to the following values: 1 – never, 2 – a few times, 3 – sometimes, 4 – often or 5 – always. Thus, items are rated from one to five according to the frequency with which they occurred, except for the items describing a pleasant childhood (2, 5, 7, 13, 19, 26, and 28), for which the

rating is inverted. The instrument assesses exposure to five types of abuse: emotional abuse (3, 8, 14, 18, 25), physical abuse (9, 11, 12, 15, and 17), sexual abuse (20, 21, 23, 24, and 27), physical neglect (1, 2, 4, 6 and 26) and emotional neglect (5, 7, 13, 19 and 28).

The CTQ-SF has good internal consistency with Cronbach's alpha values of .92 for the sexual abuse subscale (.825 our sample), .91 (.874 our sample) for emotional neglect, .87 (.844 our sample) for emotional abuse, .83 for physical abuse (.811 our sample), and .61 for physical neglect (.589 our sample) (Dias et al., 2013).

### **Attachment**

The Experiences in Close Relationships Questionnaire (ECR) consists of 36 items that assess two dimensions of adult attachment style – avoidance (18 items) and anxiety (18 items), with odd numbers corresponding to avoidance and equals to the concern scale. This questionnaire exists in a female and a male version and the answers are given on a seven-point scale (1 – Disagree strongly, 2, 3, 4 – neutral/mixed, 5, 6, and 7 – Agree strongly) with only the items marked with the extremes and the central point. Items 3, 15, 19, 22, 25, 27, 29, 31, 33 and 35 are inverted items. Higher concordance scores indicate lower levels of avoidance and concern in romantic relationships, ranging from a maximum of 252 to a minimum of 36.

The data for the Portuguese population show a high level of precision due to internal consistency, with Cronbach's alpha coefficients of .93 (.86 our sample) for the avoidance scale and .87(.85 our sample) for the concern scale (Moreira et al., 2006).

### **Method**

This study received approval by the Ethics Committee (name blinded). Participants were recruited through social networks and filled out an online survey using the platform LimeSurvey. To complete the questionnaires, participants were required to read the consent form, provide informed consent, and indicate that they were aware of the purpose and implications of their participation.

The study was conducted by all guidelines of the Declaration of Helsinki and general data protection regulations. The final anonymized database was transferred for statistical analysis using SPSS – Statistical Package for Social Sciences v. 26.0 (IBM-SPSS Inc-USA).

A Portuguese community sample was recruited via an online survey. Participant recruitment for this survey was completed using a non-list-based, non-probability sample. The essential sampling design was to spread the recruitment as broadly as possible, which requires several recruitment



**Table 2.** Correlation Table.

Measure	<i>M</i>	<i>SD</i>	2	3	4	5	6	7	8
1. Emotional Abuse	7.53	0.139	,719**	,240**	,556**	,511**	,361**	,273**	,229**
2. Emotional Neglect	8.50	0.160		,132**	,503**	,568**	,385**	,226**	,145**
3. Sexual Abuse	5.38	0.060			,179**	,199**	,167**	,099*	0,40
4. Physical Abuse	5.53	0.068				,473**	,210**	,142**	,119**
5. Physical Neglect	5.94	0.082					,311**	,232**	,188**
6. Avoidance	40.66	0.847						,345**	,189**
7. Anxiety	67.68	0.809							,282**
8. Over-responsivity									

Note: \*\*. Correlation is significant at the 0.01 level (2-tailed).

\*Correlation is significant at the 0.05 level (2-tailed).

channels to be used. Thus, subjects were recruited using advertisements on social media (mainly Facebook), personal contacts, and mailing lists.

## Results

### *Initial examination of data*

Preliminary analyses revealed that all variables in the model were significantly correlated with one another (Table 2), except sexual abuse and over-responsivity. Ten separate mediation regression analyses utilizing the PROCESS macro (model 4; Hayes, 2013) were conducted to examine whether SOR mediated the relation between the five types of child abuse (emotional abuse, emotional neglect, sexual abuse, physical abuse, physical neglect) and adult-attachment anxiety and avoidance. All five types of child abuse were entered into each regression analysis (i.e., one predictor variable controlling for the other two), attachment-related anxiety and avoidance were examined separately according to current standards within the literature (Hayes, 2009); see Figure 1 for conceptual model.

Before proceeding with the final model, several assumptions were tested. Regarding the normally distributed residuals assumption, standardized residuals displayed an approximately normal distribution for both dependent variables (skewness = 0.067; kurtosis = 0.195 for adult attachment – anxiety and skewness = 0.797; kurtosis = 0.677 for adult attachment – avoidance). One participant presented standardized residuals greater than |4.0|, suggesting that the regression equation did not accurately predict the dependent variable for these cases. Thus, these subjects were excluded from the analysis. The independence of errors assumption was not violated (Durbin–Watson statistic = 1.959 and Durbin–Watson statistic = 1.716, respectively). Finally, the scatter plot with absolute standardized residuals by standardized predicted values suggested some degree of heteroscedasticity for the adult attachment anxiety, which was further validated by the Breusch – Pagan test. Thus, the regression model was run using the wild bootstrap, which does not assume homoscedasticity (Flachaire, 2005).



Three of the categories related to the Childhood Trauma Questionnaire were over the limit for skewness and kurtosis (sexual abuse, physical abuse, and physical neglect). As such, the Hayes PROCESS Macro (Hayes, 2013) was utilized to examine the indirect effects of the mediation models through bootstrapping procedures. There were no missing values, as all the questions were necessary to proceed with the questionnaire.

**Table 3.** Model expressing mediation role of over-responsivity in emotional abuse (y=attachment-related anxiety).

R	R <sup>2</sup>	F	Df1, Df2	p
.23	.05	27.65	1.498	.000
	$\beta$	SE	t	p
<b>Constant</b>	7.73	.09	8.56	.000
<b>Emotional Abuse</b>	.58	.11	5.26	.000
Outcome: attachment-related anxiety				
.35	.13	35.63	2.497	.000
	$\beta$	se	t	p
<b>Constant</b>	51.21	2.13	24.03	.000
<b>Emotional Abuse</b>	1.28	.25	5.09	.000
<b>Over-responsivity</b>	.53	.10	5.39	.000
Direct effect of X on Y(c path)				
	Effect	SE	LLCI	ULCI
<b>Emotional Abuse</b>	1.28	.25	.78	1.77
Indirect effect of X on Y(ab path)				
<b>Over-responsivity</b>	.31	.09	.15	.51

**Table 4.** Model expressing mediation role of over-responsivity in emotional neglect (y=attachment-related anxiety).

R	R <sup>2</sup>	F	Df1, Df2	p
.15	.02	10.77	1.498	.000
	$\beta$	se	t	p
<b>Constant</b>	9.38	.90	10.40	.000
<b>Emotional Neglect</b>	.32	.10	3.28	.001
Outcome: attachment-related avoidance				
.34	.11	32.25	2.497	.000
	$\beta$	SE	t	p
<b>Constant</b>	52.05	2.17	23.96	.000
<b>Emotional Neglect</b>	.96	.22	4.44	.000
<b>Over-responsivity</b>	.58	.10	5.98	.000
Direct effect of X on Y(c path)				
	Effect	SE	LLCI	ULCI
<b>Emotional Neglect</b>	.19	.08	.05	.35
Indirect effect of X on Y(ab path)				
<b>Over-responsivity</b>	.19	.08	.05	.35

## Mediation models

### *Attachment-related anxiety*

Five separate analyses were conducted examining whether over-responsivity mediated the relation between the five types of child abuse and attachment-related anxiety. Within all five models, four of them significantly predicted the mediator, the outcome variable, and the outcome variable when the mediator was included in the model. This indicates that child abuse significantly influences adult attachment anxiety, while mediated by over-responsivity.

Regarding attachment-related anxiety, the two models with the highest  $\beta$  were physical abuse and physical neglect. For either adult attachment variable, we will report the two most significant models. The remaining models are expressed in [Tables 3 and 4](#). Physical Abuse significantly predicted the mediator ( $F(1,498) = 7.19, p = <.01, R^2 = 0.01$ ;  $\beta = 0.62, t(498) = 3.68, p = <.01$ ), the outcome, ( $F(2,497) = 25.08, p = <.01, R^2 = 0.09$ ;  $\beta = 1.29, t(497) = 2.55, p = <.01$ ), and the outcome variables when the mediator was included in the model ( $F(1,498) = 10.23, p = <.01, R^2 = 0.02$ ;  $b = 1.68, t(498) = 5.20, p = <.01$ ). This indicates that physical abuse shown statistically significant influence in attachment-related anxiety.

This model as the main predictor revealed both direct (Effect = 1.30, SE = .5097, LLCI = .0111, ULCI = .2977) and indirect (Effect = .0321, Boot SE = .0178, Boot LLCI = .0007, Boot ULCI = .0703) effects, as bias corrected bootstrap 95% confidence intervals based on 5000 bootstrap samples did not include zero: providing further evidence of mediation.

Physical neglect significantly predicted the mediator ( $F(1,498) = 18.26, p = <.01, R^2 = 0.04$ ;  $\beta = 0.81, t(498) = 4.27, p = <.01$ ), the outcome ( $F(2,497) = 41.16, p = <.01, R^2 = 0.14$ ;  $\beta = 2.04, t(497) = 7.85, p = <.01$ ), and the outcome variables when the mediator was included in the model ( $F(1,498) = 28.25, p = <.01, R^2 = 0.05$ ;  $\beta = 2.28, t(498) = 5.31, p = <.01$ ). This indicates that physical neglect shown statistically significant influence in attachment-related anxiety. This model as the main predictor revealed both direct (Effect = 1.83, SE = .4243, LLCI = .9920, ULCI = 2.656) and indirect (Effect = .4594, Boot SE = .1577, Boot LLCI = .1727, Boot ULCI = .7811) effects, as bias corrected bootstrap 95% confidence intervals based on 5000 bootstrap samples did not include zero: providing further evidence of mediation. As mentioned above, the remaining models: emotional abuse and emotional neglect, have an indirect effect of 19.5% and 16.4%, respectively.

Sexual abuse was the model that did not significantly predict the mediator ( $F(1,498) = .79, p = .38, R^2 = 0.00$ ;  $\beta = 0.234, t(498) = .89, p = .38$ ). It significantly predicted the outcome ( $F(2,497) = 23.84, p = <.01, R^2 = 0.09$ ;  $\beta = 1.191, t(497) = 2.05, p = .04$ ) and the overall model when the mediator was included ( $F(1,498) = 4.94, p = .027, R^2 = 0.01$ ;  $\beta = 1.34, t(498) = 2.22, p = .027$ ). The model with Sexual abuse has the main predictor revealed direct (Effect = 1.1915, SE = .5806, LLCI = .0407, ULCI = 2.3322) effects, but could not reveal

**Table 5.** Model expressing medication role of over-responsivity in emotional abuse (y=attachment-related avoidance).

R	R <sup>2</sup>	F	Df1, Df2	p
.23	.05	27.65	1.498	.000
	$\beta$	SE	t	p
<b>Constant</b>	7.73	.09	8.56	.000
<b>Emotional Abuse</b>	.58	.11	5.26	.000
Outcome: attachment-related avoidance				
.38	.14	41.16	2.497	.000
	$\beta$	SE	t	p
<b>Constant</b>	22.03	2.21	9.97	.000
<b>Emotional Abuse</b>	2.04	.26	7.85	.000
<b>Over-responsivity</b>	.27	.10	2.62	.000
Direct effect of X on Y(c path)				
	Effect	SE	LLCI	ULCI
<b>Emotional Abuse</b>	2.04	.26	.03	.33
Indirect effect of X on Y(ab path)				
<b>Over-responsivity</b>	.03	.01	.00	.05

**Table 6.** Model expressing medication role of over-responsivity in emotional neglect (y=attachment-related avoidance).

R	R <sup>2</sup>	F	Df1, Df2	p
.14	.02	10.77	1.498	.001
	$\beta$	se	t	p
<b>Constant</b>	9.38	.09	10.40	.000
<b>Emotional Neglect</b>	.32	.10	3.28	.001
Outcome: attachment-related avoidance				
.41	.17	49.64	2.497	.000
	$\beta$	SE	t	p
<b>Constant</b>	20.27	2.21	9.19	.000
<b>Emotional Neglect</b>	1.94	.22	8.83	.000
<b>Over-responsivity</b>	.33	.10	3.28	.001
Direct effect of X on Y(c path)				
	Effect	SE	LLCI	ULCI
<b>Emotional Neglect</b>	1.94	.22	1.50	2.37
Indirect effect of X on Y(ab path)				
<b>Over-responsivity</b>	.10	.06	.02	.24

indirect effects (Effect = .1498, Boot SE =.1460, Boot LLCI = -.1241, Boot ULCI =.54585), as the 95% confidence intervals included zero.

### ***Attachment-related avoidance***

An additional five separate analyses were conducted examining whether over-responsivity mediated the relation between the five types of child abuse and attachment-related avoidance. Within all five models, again four of them significantly predicted the mediator, the outcome variable, and the outcome variable when the mediator was included in the model. This indicates that

child abuse significantly influences adult attachment avoidance, while mediated by over-responsivity.

Regarding attachment-related avoidance, again, the two models with the highest  $\beta$  were physical abuse and physical neglect. The remaining models are expressed in [Tables 5 and 6](#), expressing an indirect effect of 7.11% and 5.12% for emotional abuse and emotional neglect respectively.

Once again, physical abuse significantly predicted the mediator ( $F(1,498) = 7.19, p = <.01, R^2 = 0.01$ ;  $\beta = 0.62, t(498) = 2.68, p = <.01$ ), the outcome ( $F(2,497) = 19.06, p = <.01, R^2 = 0.07$ ;  $\beta = 2.35, t(497) = 4.36, p = <.01$ ), and the outcome variables when the mediator was included in the model ( $F(1,498) = 22.91, p = <.01, R^2 = 0.04$ ;  $b = 2.60, t(498) = 4.79, p = <.01$ ). This indicates that physical abuse shown statistically significant influence in attachment-related avoidance. The model with physical abuse as the main predictor revealed both direct (Effect = 2.35, SE = .5393, LLCI = 1.2923, ULCI = 3.4113) and indirect (Effect = .2456, Boot SE = .1608, Boot LLCI = .0062, Boot ULCI = .6232) effects, as bias corrected bootstrap 95% confidence intervals based on 5000 bootstrap samples did not include zero: providing further evidence of mediation.

Physical neglect also significantly predicted the mediator ( $F(1,498) = 18.26, p = <.01, R^2 = 0.04$ ;  $\beta = 0.8090, t(498) = 4.27, p = <.01$ ), the outcome ( $F(2,497) = 32.01, p = <.01, R^2 = .11$ ;  $\beta = 2.94, t(497) = 6.63, p = <.01$ ), and the outcome variables when the mediator was included in the model ( $F(1,498) = 53.17, p = <.01, R^2 = 0.09$ ;  $\beta = 3.20, t(498) = 7.92, p = <.01$ ). This indicates that physical neglect shown statistically significant influence in attachment-related avoidance. This model as the main predictor revealed both direct (Effect = 2.94, SE = .4436, LLCI = 2.0713, ULCI = 3.8146) and indirect (Effect = .2626, Boot SE = .1240, Boot LLCI = .0608, Boot ULCI = .5413) effects, as bias corrected bootstrap 95% confidence intervals based on 5000 bootstrap samples did not include zero: providing further evidence of mediation.

Sexual abuse was the model that did not predict significantly the mediator ( $F(1,498) = .79, p = .38, R^2 = 0.00$ ;  $\beta = 0.234, t(498) = .89, p = .38$ ). It significantly predicted the outcome ( $F(2,497) = 16.1756, p = <.01, R^2 = 0.06$ ;  $\beta = 2.2612, t(497) = 3.6692, p = <.01$ ) but, did not predict the overall model when the mediator was included ( $F(1,498) = 14.2601, p = .38, R^2 = 0.03$ ;  $\beta = 2.3638, t(498) = 3.7763, p = <.01$ ). The model with sexual abuse as the main predictor revealed direct (Effect = 2.2612, SE = .6163, LLCI = 1.0504, ULCI = 3.4721) effects, but could not reveal indirect effects (Effect = .1026, Boot SE = .1055, Boot LLCI = -.0871, Boot ULCI = .3342), as the 95% confidence intervals included zero.

It is important to notice that although child abuse variables were significant predictors in every model, with the exception of sexual abuse, for the distinct outcome variables, there were very different mediations and effects. These important differences had additional evidence that these two constructs should be examined separately.

## Discussion

The results emphasize the importance of investigating SOR and attachment-related anxiety and avoidance when examining the impact of different types of child maltreatment. Furthermore, these findings are consistent with previous literature linking SOR and attachment-related anxiety and avoidance (Jerome & Liss, 2005), as well as child abuse to SOR (Koomar, 2009). It is important to note that the proposed model expresses higher mediation for attachment-related anxiety rather than avoidance. Generally, there is supporting evidence for a relationship between SOR and anxiety (Kinnealey & Fuiiek, 1999).

SOR is considered a condition that predisposes the individual to be hyper-vigilant for stimuli, even those that are mostly neutral, which can increase the sensation of threat. Individuals that are systematically confronted with unpleasant environmental stimuli, anxiety can become generalized and assumed as acceptable (Van Hulle et al., 2018). Research suggests that people who exhibit high sensory reactivity are more likely to experience anxiety in relationships than those with low to normal reactivity. It is plausible that those who are sensitive to sensory stimuli may be oversensitive to environmental factors in general and may naturally be more likely to worry about details in relationships that would not concern the average individual (Jerome & Liss, 2005).

These findings also emphasize the need to examine attachment-related anxiety and avoidance as separate constructs (Espeleta et al., 2017). We found that physical abuse and physical neglect are the most significant and with the highest percentage of mediation, both in attachment-related anxiety and avoidance. Another study reported that physical abuse was significantly related to the avoidant attachment style (McCarthy & Taylor, 1999). Physically abused children are described as less compliant and more impulsive than neglected children, with acting-out behavior and lack of concern or empathy toward the distress of peers and appears to become more problematic with age (Wodarski et al., 1990).

Studies have shown that preschoolers who experience neglect tend to exhibit characteristics such as social withdrawal, increased vulnerability to victimization, heightened anxiety, and decreased popularity, while also demonstrating lower levels of social competence and greater dependence on others. However, as these children grow up and enter school, some of these negative effects may become less pronounced (Schilling & Christian, 2014). Another study found that 50% of the neglected infants belong to the anxious/ambivalent group (Finzi et al., 2000).

Of the five types of child abuse severity, sexual abuse was the only maltreatment variable in the model that is not correlated to SOR. These results are consistent with a study that also found no relationship between the prevalence of atypical sensory processing behaviors and postnatal risks such as sexual abuse (Jirikowic et al., 2020). This result can be explained by the fact that in

this sample only 19 individuals were separated from the average score of 5 in sexual abuse. As this is a community sample, it is expected that most people do not suffer from sexual abuse. These results could be different if this relationship were studied in people who have been sexually abused (Jeon & Bae, 2022). Much of adult attachment and intimacy is highly dependent on how we manage sensory stimuli, for which SOR may be challenging. Individuals with sensory modulation dysfunction can be over-responsive to the sensations of touch, movement, sight, sound, and smell, thus negatively impacting their attachment to romantic partners.

### Considerations for intervention

The current results have implications for individual treatment approaches for child abuse victims. Trauma often first manifests on a somatosensory level (Gupta, 2013), causing feelings of numbness, detachment, or dissociation from the body, which can lead to difficulties processing and integrating sensory information. Incorporating sensory modulation-based interventions into established verbal-based therapies such as trauma-focused CBT or DBT may be a suitable treatment approach for adult trauma survivors. This may be particularly useful in cases where overwhelming stress impedes verbal expression of emotions.

Therapy approaches that utilize bottom-up and body-centered techniques have the potential to be an effective treatment option for trauma survivors (Van der Kolk, 2014). These approaches focus on body responses to sensory input rather than thought processes when addressing emotional disorders. There is evidence from case studies that utilizing sensory inputs (e.g., vestibular, proprioceptive, and tactile) can effectively reduce behavior problems and symptoms of post-traumatic stress in children and adolescents (Warner et al., 2014).

### Limitations

While these novel findings contribute to our understanding of the relationship between child abuse severity, SOR, and attachment-related anxiety and avoidance within intimate relationships, we must temper our interpretation of findings by considering the limitations of the study. First, data collection was conducted using an online survey which ultimately does not allow to represent the whole population. Given the online limitation, we had a gender class imbalance, being that male participants were only 29.4% of our sample, skewing the results regarding gender.

Some potential participants do not have access to the Internet or were not exposed to our online recruitment/advertising endeavors, not allowing them to be recruited for this study. The questionnaire was fairly extensive which might have made the answers less credible due to the participants being bored.

Finally, this study utilized retrospective, self-report data thereby limiting the ability to infer temporal or causal relations. Additional longitudinal studies would be valuable to understand how SOR, child maltreatment, and attachment develop over time.

Although other studies have examined relationships among two of the three variables studied here (SOR, child abuse, and attachment-related anxiety and avoidance), this is the first study the authors are aware of that examines all variables in one model. The inclusion of SOR within this model provides a clear point of intervention for child abuse survivors experiencing attachment difficulties within romantic relationships. Therefore, it should be considered to address sensory issues and SOR in counseling and occupational therapy interventions to approach problems in romantic relationships that are related to attachment-related anxiety or avoidance. Researchers have long sought to understand how experiences in childhood are interpreted and translated into behaviors and interpersonal styles in adulthood.

## Conclusion

This study is one piece of the puzzle that attempts to unravel the relation between child abuse experiences and attachment-related anxiety and avoidance in adult intimate relationships. The current study suggests that SOR expresses higher mediation for attachment-related anxiety rather than avoidance. By delving deeper into the underlines of the specific mechanisms connecting these variables, we can better target our interventions to address the specific needs of individuals with SOR and help them learn and implement strategies to manage their social interactions. Additionally, it is important to note that physical abuse and neglect presented the highest mediation in both attachment-related anxiety and avoidance.

Understanding the impact of child abuse and neglect on attachment and interpersonal functioning is essential to providing effective interventions to survivors. Tailoring interventions to the specific needs of each individual can help them overcome the effects of past trauma and develop resilience for the future. Our research opens up the possibility that incorporating sensory modulation-based interventions into trauma-focused therapies may contribute to improving attachment and adult intimate relationships among survivors. Future research is required to determine the effectiveness of this approach for individuals with different types of trauma experiences.

Through this enlightened knowledge regarding SOR, attachment-related anxiety and avoidance, we can improve our ability to assist survivors in growing and remaining resilient throughout adulthood, and ultimately help



them strive in their lives, including relationships goals and professional achievements.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## ORCID

Simão Ferreira  <http://orcid.org/0000-0001-8233-2217>

Susana Aguiar Rocha  <http://orcid.org/0000-0002-5833-165X>

Nuno Barbosa-Rocha  <http://orcid.org/0000-0002-3139-2786>

## References

- APAV, A. P. d. A. à. V. (2021) . *Relatório Anual 2020*. [WWW.APAV.PT/ESTATISTICAS](http://WWW.APAV.PT/ESTATISTICAS) .
- Arata, C. M., Langhinrichsen-Rohling, J., Bowers, D., & O'farrill-Swails, L. (2005). Single versus multi-type maltreatment. *Journal of Aggression, Maltreatment & Trauma*, 11(4), 29–52. [https://doi.org/10.1300/J146v11n04\\_02](https://doi.org/10.1300/J146v11n04_02)
- Bacon, H., & Richardson, S. (2001). Attachment theory and child abuse: An overview of the literature for practitioners. *Child Abuse Review*, 10(6), 377–397. <https://doi.org/10.1002/car.718>
- Bar-Shalita, T., & Cermak, S. A. (2020). Multi-sensory responsiveness and personality traits predict daily pain sensitivity [original research]. *Frontiers in Integrative Neuroscience*, 13 (77). <https://doi.org/10.3389/fnint.2019.00077>
- Bolger, K. E., Patterson, C. J., & Kupersmidt, J. B. (1998). Peer relationships and self-esteem among children who have been maltreated. *Child Development*, 69(4), 1171–1197. <https://doi.org/10.1111/j.1467-8624.1998.tb06166.x>
- Bowlby, J. (1982). Attachment and loss: Retrospect and prospect. *The American Journal of Orthopsychiatry*, 52(4), 664–678. <https://doi.org/10.1111/j.1939-0025.1982.tb01456.x>
- Bundy, A. C., & Lane, S. J. (2019). *Sensory integration: Theory and practice*. F.A. Davis Company.
- Christ, C., de Waal, M. M., Dekker, J. J. M., van Kuijk, I., van Schaik, D. J. F., Kikkert, M. J., Goudriaan, A. E., Beekman, A. T. F., Messman-Moore, T. L., & Seedat, S. (2019). Linking childhood emotional abuse and depressive symptoms: The role of emotion dysregulation and interpersonal problems. *PLoS One*, 14(2), e0211882. <https://doi.org/10.1371/journal.pone.0211882>
- Cohen, J. R., Menon, S. V., Shorey, R. C., Le, V. D., & Temple, J. R. (2017). The distal consequences of physical and emotional neglect in emerging adults: A person-centered, multi-wave, longitudinal study. *Child Abuse & Neglect*, 63, 151–161. <https://doi.org/10.1016/j.chiabu.2016.11.030>
- Dias, A., Sales, L., Carvalho, A., Vale, I. C., Kleber, R., & Cardoso, R. M. (2013). Estudo de propriedades psicométricas do Questionário de Trauma de Infância – Versão breve numa amostra portuguesa não clínica.
- Dunn, W. (2007). Supporting children to participate successfully in everyday life by using sensory processing knowledge. *Infants & Young Children*, 20(2), 84–101. <https://doi.org/10.1097/01.IYC.0000264477.05076.5d>
- Espeleta, H. C., Palasciano-Barton, S., & Messman-Moore, T. L. (2017). The impact of child abuse severity on adult attachment anxiety and avoidance in college women: The role of

- emotion dysregulation. *Journal of Family Violence*, 32(4), 399–407. <https://doi.org/10.1007/s10896-016-9816-0>
- Finzi, R., Cohen, O., Sapir, Y., & Weizman, A. (2000). Attachment styles in maltreated children: A comparative study. *Child Psychiatry and Human Development*, 31(2), 113–128. <https://doi.org/10.1023/A:1001944509409>
- Flachaire, E. (2005). Bootstrapping heteroskedastic regression models: Wild bootstrap vs. pairs bootstrap. *Computational Statistics & Data Analysis*, 49(2), 361–376. <https://doi.org/10.1016/j.csda.2004.05.018>
- Greenberg, D. M., Baron-Cohen, S., Rosenberg, N., Fonagy, P., Rentfrow, P. J., & Matosin, N. (2018). Elevated empathy in adults following childhood trauma. *PLoS One*, 13(10), e0203886. <https://doi.org/10.1371/journal.pone.0203886>
- Gubbels, J., Assink, M., Prinzie, P., & van der Put, C. E. (2021). Why healthcare and education professionals underreport suspicions of child abuse: A qualitative study. *Social Sciences*, 10(3), 98. <https://doi.org/10.3390/socsci10030098>
- Gupta, M. A. (2013). Review of somatic symptoms in post-traumatic stress disorder. *International Review of Psychiatry*, 25(1), 86–99. <https://doi.org/10.3109/09540261.2012.736367>
- Hayes, A. F. (2009). Beyond baron and Kenny: Statistical mediation analysis in thenewmillennium. *Communication Monographs*, 76(4), 408–420. <https://doi.org/10.1080/03637750903310360>
- Hayes, A. F. (2013). *Introduction to mediation, moderation, and conditional process analysis: A regression-based approach*. Guilford Press.
- Jedd, K., Hunt, R. H., Cicchetti, D., Hunt, E., Cowell, R. A., Rogosch, F. A., Toth, S. L., & Thomas, K. M. (2015). Long-term consequences of childhood maltreatment: Altered amygdala functional connectivity. *Development and Psychopathology*, 27(4 Pt 2), 1577–1589. <https://doi.org/10.1017/S0954579415000954>
- Jeon, M. S., & Bae, E. B. (2022). Emotions and sensory processing in adolescents: The effect of childhood traumatic experiences. *Journal of Psychiatric Research*, 151, 136–143. <https://doi.org/10.1016/j.jpsychires.2022.03.054>
- Jerome, E. M., & Liss, M. (2005). Relationships between sensory processing style, adult attachment, and coping. *Personality and Individual Differences*, 38(6), 1341–1352. <https://doi.org/10.1016/j.paid.2004.08.016>
- Jirikowic, T. L., Thorne, J. C., McLaughlin, S. A., Waddington, T., Lee, A. K. C., & Astley Hemingway, S. J. (2020). Prevalence and patterns of sensory processing behaviors in a large clinical sample of children with prenatal alcohol exposure. *Research in Developmental Disabilities*, 100, 103617. <https://doi.org/10.1016/j.ridd.2020.103617>
- Joseph, R., Casteleijn, D., Linde, J., & Franzsen, D. (2021). Sensory modulation dysfunction in child victims of trauma: A scoping review. *Journal of Child & Adolescent Trauma*, 14(4), 455–470. <https://doi.org/10.1007/s40653-020-00333-x>
- Kinnealey, M., & Fuiek, M. (1999). The relationship between sensory defensiveness, anxiety, depression and perception of pain in adults. *Occupational Therapy International*, 6(3), 195–206. <https://doi.org/10.1002/oti.97>
- Koomar, J. A. (2009). Trauma-and attachment-informed sensory integration assessment and intervention. *Sensory Integration Special Interest Section Quarterly*, 32(4), 1–4.
- Maier, A., Gieling, C., Heinen-Ludwig, L., Stefan, V., Schultz, J., Güntürkün, O., Becker, B., Hurlemann, R., & Scheele, D. (2020). Association of childhood maltreatment with interpersonal distance and social touch preferences in adulthood. *The American Journal of Psychiatry*, 177(1), 37–46. <https://doi.org/10.1176/appi.ajp.2019.19020212>
- Maier, A., Heinen-Ludwig, L., Güntürkün, O., Hurlemann, R., & Scheele, D. (2020). Childhood maltreatment alters the neural processing of chemosensory stress signals. *Frontiers in Psychiatry*, 11, 783. <https://doi.org/10.3389/fpsy.2020.00783>

- McCarthy, G., & Taylor, A. (1999). Avoidant/Ambivalent attachment style as a mediator between abusive childhood experiences and adult relationship difficulties. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 40(3), 465–477. <https://doi.org/10.1111/1469-7610.00463>
- McGlone, F., Wessberg, J., & Olausson, H. (2014). Discriminative and affective touch: Sensing and feeling. *Neuron*, 82(4), 737–755. <https://doi.org/10.1016/j.neuron.2014.05.001>
- McGreevy, S., & Boland, P. (2020). Sensory-based interventions with adult and adolescent trauma survivors. *Irish Journal of Occupational Therapy*, 48(1), 31–54. <https://doi.org/10.1108/IJOT-10-2019-0014>
- Miller, L. J., Nielsen, D. M., & Schoen, S. A. (2012). Attention deficit hyperactivity disorder and sensory modulation disorder: A comparison of behavior and physiology. *Research in Developmental Disabilities*, 33(3), 804–818. <https://doi.org/10.1016/j.ridd.2011.12.005>
- Moreira, J. M., Lind, W., Santos, M. J., Moreira, A. R., Gomes, M. J., Justo, J., Oliveira, A. P., Filipe, L. A., & Faustino, M. (2006). “Experiências em Relações Próximas”, um questionário de avaliação das dimensões básicas dos estilos de vinculação nos adultos: Tradução e validação para a população Portuguesa.
- Mulligan, S., Schoen, S., Miller, L., Valdez, A., Wiggins, A., Hartford, B., & Rixon, A. (2019). Initial Studies of validity of the sensory processing 3-dimensions Scale. *Physical & Occupational Therapy in Pediatrics*, 39(1), 94–106. <https://doi.org/10.1080/01942638.2018.1434717>
- Rueness, J., Myhre Md, M. C., Strøm, I. F., Wentzel Larsen, T., Dyb, G., & Thoresen, S. (2020). Child abuse and physical health: A population-based study on physical health complaints among adolescents and young adults. *Scandinavian Journal of Public Health*, 48(5), 511–518. <https://doi.org/10.1177/1403494819848581>
- Schilling, S., & Christian, C. W. (2014). Child physical abuse and neglect. *Child and Adolescent Psychiatric Clinics of North America*, 23(2), 309–319, ix. <https://doi.org/10.1016/j.chc.2014.01.001>
- Schoen, S. A., Miller, L. J., & Sullivan, J. (2017). The development and psychometric properties of the sensory processing scale inventory: A report measure of sensory modulation. *Journal of Intellectual & Developmental Disability*, 42(1), 12–21. <https://doi.org/10.3109/13668250.2016.1195490>
- Stoltenborgh, M., Bakermans-Kranenburg, M. J., Alink, L. R. A., & van IJzendoorn, M. H. (2015). The prevalence of child maltreatment across the globe: Review of a series of meta-analyses. *Child Abuse Review*, 24(1), 37–50. <https://doi.org/10.1002/car.2353>
- Unger, J. A. M., & De Luca, R. V. (2014). The relationship between childhood physical abuse and adult attachment styles. *Journal of Family Violence*, 29(3), 223–234. <https://doi.org/10.1007/s10896-014-9588-3>
- Van der Kolk, B. (2014). *The body keeps the score: Mind, brain and body in the transformation of trauma*. penguin UK.
- Van Hulle, C. A., Lemery-Chalfant, K., & Hill Goldsmith, H. (2018). Parent-offspring transmission of internalizing and sensory over-responsivity symptoms in adolescence. *Journal of Abnormal Child Psychology*, 46(3), 557–567. <https://doi.org/10.1007/s10802-017-0300-y>
- Warner, E., Spinazzola, J., Westcott, A., Gunn, C., & Hodgdon, H. (2014). The body can change the score: Empirical support for somatic regulation in the treatment of traumatized adolescents. *Journal of Child & Adolescent Trauma*, 7(4), 237–246. <https://doi.org/10.1007/s40653-014-0030-z>
- Whiting, C. C. (2018). Trauma and the role of the school-based occupational therapist. *Journal of Occupational Therapy, Schools, & Early Intervention*, 11(3), 291–301. <https://doi.org/10.1080/19411243.2018.1438327>
- WHO. (1999). *Report of the consultation on child abuse prevention*. World Health Organization.
- WHO. (2022). *Violence Info - Child maltreatment*. World Health Organization.

- Witte, S., Fegert, J. M., & Walper, S. (2020). Sibling relationship pattern in the context of abuse and neglect: Results from a sample of adult siblings. *Child Abuse & Neglect*, 106, 104528. <https://doi.org/10.1016/j.chiabu.2020.104528>
- Wodarski, J. S., Kurtz, P. D., Gaudin, J. M., Jr., & Howing, P. T. (1990). Maltreatment and the school-age child: Major academic, socioemotional, and adaptive outcomes. *Social Work*, 35 (6), 506–513. <https://doi.org/10.1093/sw/35.6.506>
- Yoon, D. (2020). Peer-relationship patterns and their association with types of child abuse and adolescent risk behaviors among youth at-risk of maltreatment. *Journal of Adolescence*, 80 (1), 125–135. <https://doi.org/10.1016/j.adolescence.2020.02.008>