

Oral Communication 6

## Enantiomeric biodistribution and toxicity of 3-chloromethcathinone (3-CMC) in Wistar rats after acute exposure – preliminary data

**Ivan Langa**<sup>1,2,3,4,\*</sup>, **Carolina Rocha-Pereira**<sup>1,2</sup>, **Nuno Milhazes**<sup>1,2</sup>, **Diana Dias da Silva**<sup>1,3,4,5,6</sup>, **Susana Domingues**<sup>1,2</sup>, **Paula Silva**<sup>7</sup>, **Joana Barbosa**<sup>1,2</sup>, **Juliana Faria**<sup>1,2</sup>, **Maria Elizabeth Tiritan**<sup>1,2,8,9</sup> and **Cláudia Ribeiro**<sup>1,2</sup>

<sup>1</sup> Associate Laboratory i4HB – Institute for Health and Bioeconomy, University Institute of Health Sciences – CESPU, 4585-116 Gandra, Portugal

<sup>2</sup> UCIBIO – Applied Molecular Biosciences Unit, Translational Toxicology Research Laboratory, University Institute of Health Sciences (1H-TOXRUN, IUCS-CESPU), 4585-116 Gandra, Portugal

<sup>3</sup> UCIBIO – Applied Molecular Biosciences Unit, Laboratory of Toxicology, Faculty of Pharmacy, University of Porto, 4050-313 Porto, Portugal

<sup>4</sup> Associate Laboratory i4HB – Institute for Health and Bioeconomy, University of Porto, 4050-313 Porto, Portugal

<sup>5</sup> REQUIMTE/LAQV, ESS, Polytechnic of Porto, Rua Dr. António Bernardino de Almeida, 400 4200-072, Porto

<sup>6</sup> UCIBIO – Applied Molecular Biosciences Unit, Forensics and Biomedical Sciences Research Laboratory, University Institute of Health Sciences (1H-TOXRUN, IUCS-CESPU), 4585-116 Gandra, Portugal

<sup>7</sup> Departamento de Microscopia, Laboratório de Histologia e Embriologia, Instituto de Ciências Biomédicas de Abel Salazar (ICBAS), Universidade do Porto, 4050-313 Porto, Portugal

<sup>8</sup> Interdisciplinary Center for Marine and Environmental Research (CIIMAR), Port of Leixões Cruise Terminal, s/n, Matosinhos, Portugal

<sup>9</sup> Laboratory of Organic and Pharmaceutical Chemistry, Department of Chemical Sciences, Faculty of Pharmacy, University of Porto, 4050-313 Porto, Portugal

\* Correspondence: [ivan.mlanga4@gmail.com](mailto:ivan.mlanga4@gmail.com)

### Abstract

**Background:** There has been a surge in global attention to New Psychoactive Substances (NPS) [1]. Synthetic cathinones stand out as a widely consumed NPS class. Notably, 3-chloromethcathinone (3-CMC) accounted for over 34% of NPS seizures in 2021 [2], which underscores concerns regarding its consumption and health effects. Of note, 3-CMC is chiral and mostly sold as a racemate. As human metabolism and pharmacological effects can be enantioselective [3], determination of the impact of enantioselectivity in toxicokinetics/toxicodynamics is essential for the assessment of 3-CMC effects. **Objective:** This work aimed to evaluate *in vivo* the enantioselective biodistribution and toxicity of racemic 3-CMC, after an acute exposure to 3-CMC. **Methods:** Ten-week-old male Wistar rats were administered intraperitoneally with saline or 3-CMC (10 or 20 mg/kg; n=6). Twenty-four hours after, animals were deeply anesthetized and nine organs (brain, liver, kidneys, lungs, heart, spleen, gut, muscle, adipose tissue), blood and urine were collected. For evaluation of the enantiomeric biodistribution, a previous *in house* established indirect method by gas chromatography [3], was adapted and validated. Some biochemical analysis was performed using an analyser, whereas TBARS, ATP, glutathione and total protein were determined by spectrophotometry. Organs were also processed for histological analysis. **Results:** After 24 h, 3-CMC was not found in most organs. Both enantiomers were detected in urine with one dominant enantiomer, suggesting enantioselectivity in metabolism. The histopathological results showed possible central chromatolysis in the brain (20 mg/kg), liver inflammation, renal lesions, lungs' haemoptysis, and alveolar haemorrhage, in most 3-CMC-exposed animals. No differences were observed in the heart. **Conclusions:** Our findings show rapid 3-CMC renal elimination, with enantioselectivity in metabolism. Although biochemical evaluations are ongoing, the results are expected to give further insights on the 3-CMC toxicity and histological abnormalities found in the brain, kidneys, liver and lungs.

**Keywords:** new psychoactive substances; drugs of abuse; enantioselectivity; toxicokinetics; risk assessment

---

### Acknowledgments

This project was financed by FCT–Portuguese Foundation for Science and Technology, through the FCT/MCTES (PIDDAC), under the project PTDC/CTA-AMB/6686/2020 (ENANTIOTOX) and partially supported by national funds by FCT within the scope of Base Funding UIDB/04423/2020 and UIDP/04423/2020, UIDP/04423/2020 (Group of Marine Natural Products and Medicinal Chemistry-CIIMAR). Ivan Langa acknowledges a Ph.D. research grant provided by FCT 2022.12874.BD.

### References

1. Lei nº 55/2023 de 8 de setembro. Diário da República. <https://data.dre.pt/eli/lei/55/2023/09/08/p/dre/pt/html>
2. EMCDDA, European Drug Report 2023: Trends and Developments. 2023.
3. Gonçalves, R.; Ribeiro, C.; Cravo, S.; Cunha, S.C.; Pereira, J.A.; Fernandes, J.; Afonso, C.; Tiritan, M.E. Multi-residue method for enantioseparation of psychoactive substances and beta blockers by gas chromatography–mass spectrometry. *Journal of Chromatography B*, 1125 (2019) 121731.



In *Scientific Letters*, works are published under a CC-BY license (Creative Commons Attribution 4.0 International License at <https://creativecommons.org/licenses/by/4.0/>), the most open license available. The users can share (copy and redistribute the material in any medium or format) and adapt (remix, transform, and build upon the material for any purpose, even commercially), as long as they give appropriate credit, provide a link to the license, and indicate if changes were made (read the full text of the license terms and conditions of use at <https://creativecommons.org/licenses/by/4.0/legalcode>).