

Results: Of the 232 subjects evaluated, 40.1% were classified as MHO (n = 93) and 59.9% as MHUO (n = 139). Age (years), weight (Kg), BMI and WC (cm) mean (\pm standard deviation) values were $39.6 \pm 11.1/43.6 \pm 10.1$, $117.5 \pm 17.1/117.8 \pm 20.0$, $42.9 \pm 4.5/42.4 \pm 4.9$ and $118.0 \pm 11.9/120.7 \pm 14.2$ for the MHO and MHUO groups, respectively. Accordingly, in these groups, circulating uric acid concentrations (mg/dL) were 5.0 ± 1.1 and 5.4 ± 1.4 ($p = 0.033$), respectively. Hyperuricemia was present in 30.1% (n = 28) and 34.5% (n = 48) of the MHO and MHUO groups (6.2 ± 0.7 and 6.8 ± 1.3 , respectively; $p = 0.042$). Among the features included in the MHUO definition, WC, TG and glucose were positively correlated with uric acid ($r = 0.275/p < 0.001$; $r = -0.199/p = 0.002$ and $r = 0.156/p = 0.017$, respectively) while HDL-c was negatively correlated ($r = -0.188/p = 0.004$). Circulating uric acid concentrations increased with the increasing number of the features of the MHUO definition ($B = 0.153$; $p = 0.045$).

Conclusion: In our Brazilian population, hyperuricemia was present in both obesity phenotypes, but higher in the MHUO. In this phenotype, as individuals had more components of the MHUO definition, the circulating uric acid levels were higher.

References:

1. Harald Mangge et al. Obesity (2013) 21: E71-E77.
2. Matthias Bluher. European Journal of Endocrinology (2014) 171: R209-R219.
3. Sérgio Lamego Rodrigues et al. Arq Bras Cardiol. (2012) 98(1):13-21.
4. Grundy SM et al. Circulation (2005) 112 (17): 2735-52.

4 – Metabolic Syndrome and Inflammation: Is There a Microvascular and an Incretin System Impairment in the Gastrointestinal Tract?

Costa J¹, Almeida J^{2,3}, Coelho P^{2,3}, Oliveira S¹, Prudêncio C^{2,3}, Soares R^{3,4}, Gomes-Guerreiro S^{2,3,4}, Fernandes R^{2,3}

¹ Mestrado em Bioquímica em Saúde, Escola Superior de Saúde, Instituto Politécnico do Porto, Porto, Portugal

² Ciências Químicas e das Biomoléculas, Centro de Investigação em Saúde e Ambiente, Escola Superior de Saúde, Instituto Politécnico do Porto, Porto, Portugal

³ I3S – Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Porto, Portugal

⁴ Departamento de Bioquímica, Faculdade de Medicina da Universidade do Porto, Porto, Portugal

Introduction: Metabolic syndrome is a multifactorial disorder characterized by increased plasma levels of glucose, cholesterol and triglycerides, but also overweight and obesity promoted by increase of body fat mass, alterations in oxidative stress, chronic low grade inflammation and resistance to insulin leading to risk of cardiovascular diseases. ⁽¹⁾ The stomach and the intestine have an essential role in metabolism with functions of digesting food and absorption of nutrients. Also, the intestine produces incretin hormones, such as GLP-1 which regulates glucose metabolism and processes of the gastrointestinal tract. ⁽²⁾

Objectives: We aim to evaluate the inflammatory status, blood and lymphatic microvasculature and the incretin system in the intestine and stomach in animals exerting metabolic syndrome.

Methods: It was assessed the expression of IL-6, 3-NT, CD31, LYVE-1 and GLP-1 receptor on tissues from the stomach and intestine of C57BL/6 mice, divided in two groups, a high fat diet and a normal diet group.

Discussion and Conclusion: The results suggest that metabolic syndrome promotes alterations in the inflammation status, vascularization and in the incretin system of gastrointestinal tract. In high fat diet mice, it was observed higher levels of inflammation, no alteration in expression of 3-NT, lower blood microvascular density, lower number of lymphatic vessels and decreased expression of GLP-1 receptor. Further studies are important to understand other molecular mechanisms involved in metabolic syndrome and their possible influence in the gastrointestinal tract.

Keywords: metabolic syndrome, inflammation, incretin system, microvasculature, gastrointestinal tract

References:

1. Feakins RM. Obesity and metabolic syndrome: pathological effects on the gastrointestinal tract. Histopathology. 2016; 68(5): 630-40.
2. Baggio LL, Drucker DJ. Biology of incretins: GLP-1 and GIP. Gastroenterology. 2007; 132(6): 2131-57.

5 – Is the MTHFR C677T Polymorphism Associated with Obesity Risk? – a Meta-Analysis.

Costa J¹, Oliveira R^{2,3}, Prudêncio C^{1,4}, Fernandes R^{1,4}

¹Mestrado em Bioquímica em Saúde, Escola Superior de Saúde, Instituto Politécnico do Porto, Porto, Portugal

²Faculdade de Medicina da Universidade do Porto, Porto, Portugal

³Bioestatística, Biomatemática e Bioinformática, Escola Superior de Saúde, Instituto Politécnico do Porto, Porto, Portugal

⁴Ciências Químicas e das Biomoléculas, Centro de Investigação em Saúde e Ambiente, Escola Superior de Saúde, Instituto Politécnico do Porto, Porto, Portugal

Background: Overweight and obesity are a major worldwide health problem and its incidence is increasing every year. ⁽¹⁾ Methylene tetrahydrofolate reductase (MTHFR) plays an important role in folate metabolism and as a regulator of DNA methylation, synthesis, and repair. MTHFR gene is polymorphic at nucleotides 677 (C→T) and 1298 (A→C). MTHFR C677T polymorphism results in alloenzymes with decreased activity and several studies have pointed to association between the MTHFR C677T polymorphism and overweight/obesity risk. ⁽²⁾

Objectives: The present study aims to contribute to the elucidation of the impact of any C677T overweight/obesity association through a meta-analysis study of published case control studies.

Material and Methods: Pubmed, Google Scholar, Elsevier and Cochrane trials databases were searched for case control studies of associations between MTHFR C677T polymorphism and overweight/obesity. Odds ratios (ORs) with 95% confidence intervals (CIs) were estimated to assess the association.

Results: Our results suggest that MTHFR C677T polymorphisms may modify the association between intracellular folate levels and overweight/obesity risk. Homozygous individuals for the MTHFR 677T polymorphism may have a significantly increase of obesity risk.

Conclusion: Our results suggest an association between MTHFR C677T polymorphism and risk of being overweight/obese.

Keywords: MTHFR, C677T, polymorphism, obesity, risk.

References:

1. Mendonca F, Soares R. Obesity and cancer phenotype: Is angiogenesis a missed link? Life Sci. 2015;139:16-23.
2. Liew SC, Gupta ED. Methylene tetrahydrofolate reductase (MTHFR) C677T polymorphism: epidemiology, metabolism and the associated diseases. Eur J Med Genet. 2015;58(1):1-10.

6 – Polyphenols Modulate Type 2 Diabetes: Relevance to Angiogenic Paradox

Costa R^{1,2*}, Rodrigues I¹, Guardão L³, Negrão, R^{1,2}, Soares R^{1,2}

¹ Department of Biochemistry, Faculty of Medicine of the University of Porto, Porto, Portugal *mrcosta@med.up.pt

² I3S, Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Portugal

³ Animal Facility, Faculty of Medicine of the University of Porto, Porto, Portugal

Background: Diabetes mellitus (DM) is responsible for metabolic de-