

**Keywords:** *Aged, Physician-Patient Relations, Geriatrics, Empathy, Aging, Communication.*

## References

- [1] Gomes M. A comunicação e relação médico-doente como benefício para o doente e para a decisão médica. *Revista Ordem dos Médicos* No148. 2014; 79–81.
- [2] Amalraj S, Starkweather C, Nguyen C, Naeim A. Health literacy, communication, and treatment decision-making in older cancer patients. *Oncology*. 2009; 23(4): 369–75.
- [3] Giampieri M. Communication and informed consent in elderly people. *Mínerva Anestesiol*. 2012; 78(2): 236–42.

## P185

### Pharmacy professionals as providers of rapid testing for HIV infection screening

Micael Gomes<sup>1</sup>, Ângelo Jesus<sup>2</sup>, Agostinho Cruz<sup>3</sup>

<sup>1</sup>CISA, Escola Superior de Saúde, Instituto Politécnico do Porto, Portugal, [micaelmg-99@hotmail.com](mailto:micaelmg-99@hotmail.com)

<sup>2</sup>CISA, Escola Superior de Saúde, Instituto Politécnico do Porto, Portugal, [acj@ess.ipp.pt](mailto:acj@ess.ipp.pt)

<sup>3</sup>CISA, Escola Superior de Saúde, Instituto Politécnico do Porto, Portugal, [agostinhocruz@ess.ipp.pt](mailto:agostinhocruz@ess.ipp.pt)

**BACKGROUND:** Portugal continues to have one of the highest HIV rates in the European Union<sup>[1]</sup>. More than 20% of the infected population is unaware of the state of their infection and may unknowingly contribute to the spread of the disease. In addition, those who are not aware that they are infected, lose the opportunity to receive treatment that prolongs life at the onset of infection, and helps to reduce HIV / AIDS-related racial / ethnic health disparities<sup>[2-5]</sup>. The inclusion of rapid HIV screening tests in community pharmacies may improve the detection of undiagnosed infections<sup>[3]</sup>. Pharmacies are valuable partners in health promotion efforts. They can be an accessible health resource for the inclusion of HIV testing, increasing the availability, convenience, and accessibility of these testing services<sup>[4]</sup>. However, concerns have been raised, by the public and health professionals, regarding the readiness of community pharmacies to provide such tests and the willingness and preparation of pharmacy professionals to act as providers of rapid testing for HIV infection screening. **OBJECTIVE:** This paper aims to assess the benefits and barriers associated with the introduction of HIV testing in pharmacies and ascertain the readiness and availability of pharmacy professionals to provide this service. **METHODS:** A questionnaire was applied to pharmacists and registered pharmacy technicians who worked in community pharmacies of the county of Braga, selected by a convenience sample. Data from 64 pharmacy professionals was retrieved. **RESULTS:** Ninety five per cent of the pharmacies were physically ready to provide this type of service, but only about 50% would be available at this point to start providing the screenings. Pharmacy professionals were well aware of the different types of tests available and were willing to act (58%) as providers of the HIV- screening. Most of the respondents consider that these screenings provide "greater control of the disease" (27.3%), "quick results" (25.6%), and accessibility" (25%). Nevertheless, 95% of respondents considered that mandatory previous training should be required, namely on "how to deal with positive results", but also on the fundamentals of HIV infection, transmission and treatment options. **CONCLUSIONS:** This exploratory study shows that pharmacies are physically ready to provide these screenings and that most professionals are willing to participate. Nevertheless, a focus on mandatory training and general rule of conduct must be set, in order to provide a quality service.

**Keywords:** *Pharmacies/organization & administration; Mass Screening/methods; HIV Infections/diagnosis*

## References:

- [1] Amesty S, Blaney S, Crawford N, Rivera A, Fuller C. Pharmacy staff characteristics associated with support for pharmacy-based HIV testing. *Journal of the American Pharmacists Association*. 2015; 52.
- [2] Amesty SD, Crawford N, Nandi V, Perez-Figueroa R, Rivera A, Sutton M, et al. Evaluation of Pharmacy-Based HIV Testing in a High-Risk New York City Community. In *AIDS Patient Care and STDs*. Mary Ann Liebert, Inc. 2015.
- [3] Ariën K, Abrahams A, Quiñones-Mateu M, Kestens L, Vanham G, Arts E. The replicative fitness of primary human immunodeficiency virus type 1 (HIV-1) group M, HIV-1 group O, and HIV-2 isolates. *Journal of Virology*. 2005; 79: 8979-8990.
- [4] Bousquet P, Daures J, Demoly P. Principles, characteristics and interpretation of diagnostic and screening tests. 2005.
- [5] Barin F, Simon F. *Virologie*. In *Les outils du dépistage de l'infection par le VIH: concepts, progrès et limites*. John Libbey Eurotext; 2013.

## P186

### Tablet Splitting: Counselling the Patients Regarding Techniques and Tablet Format

Ana Alexandre<sup>1</sup>, Ângelo Jesus<sup>2</sup>, Agostinho Cruz<sup>3</sup>

<sup>1</sup>CISA, Escola Superior de Saúde, Instituto Politécnico do Porto, Portugal, [anabalalexandre@gmail.com](mailto:anabalalexandre@gmail.com)

<sup>2</sup>CISA, Escola Superior de Saúde, Instituto Politécnico do Porto, Portugal, [acj@ess.ipp.pt](mailto:acj@ess.ipp.pt)

<sup>3</sup>CISA, Escola Superior de Saúde, Instituto Politécnico do Porto, Portugal, [agostinhocruz@ess.ipp.pt](mailto:agostinhocruz@ess.ipp.pt)

**BACKGROUND:** Tablet splitting consists in subdividing a tablet in order to facilitate its intake or to provide smaller doses, although its efficacy depends on many factors, such as tablet shape, size, splitting technique/device and patient ability<sup>[1,2]</sup>. This technique is commonly used in households. Problems associated with this technique are mainly due to the lack of ability to obtain equal parts after breaking the tablet, either due to physical characteristics or the patient's ability to do so<sup>[3]</sup>, and the loss of mass due to crumbles<sup>[4]</sup>. **OBJECTIVE:** The objectives of this study are to assess if there are any significant differences when untrained volunteers split previously selected scored tablets, by analysing the results regarding tablet format, splitting method, loss of mass and the probability of obtaining accurate halves. **METHODS:** Formulations identified by a team of pharmacy technicians as being often split in hospital practice were gathered. A total of sixty tablets were randomly chosen from each formulation and were weighted using a Kern Abs 220-4 analytical scale. The records for each formulation were then inserted in a Microsoft® Excel 2010 spreadsheet, totalling 90 results per formulation/method combination. The software was then used to verify which tablets complied with the specifications from the European Pharmacopoeia for splitting tablets and which did not have a loss of mass greater than 3%. As for splitting accuracy, the European Pharmacopoeia calls for only one half of each tablet to be weighted. For the loss of mass test, a sum of the weights of both halves was used. **RESULTS:** Regarding splitting accuracy only 40% passed the European Pharmacopoeia criteria. Regarding loss of mass, 60% presented at least one tablet with a loss of mass greater than 3%. Only 30% of the oral drug formulations complied with all the criteria established. No difference was found regarding the splitting method. **CONCLUSIONS:** Having a scoring line does not assure that it is advisable to split the tablet, since it may lead to unequal halves and/or significant loss of mass. The Pharmacy Professional should take these informations into account when counselling the patient so that he or she will obtain accurate halves and the loss of mass is not significant.

**Keywords:** *Tablets/administration & dosage;*

#### References:

- [1] Teng J, Song CK, Williams RL, Polli JE. Lack of medication dose uniformity in commonly split tablets. *J Am Pharm Assoc.* 2002; 42(2): 195-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11926662>
- [2] van Santen E, Barends DM, Frijlink HW. Breaking of scored tablets: a review. *Eur J Pharm Biopharm.* 2002; 53(2): 139-45. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11879995>
- [3] Quinzler R, Gasse C, Schneider A, Kaufmann-Kolle P, Szeceenyi J, Haefeli WE. The frequency of inappropriate tablet splitting in primary care. *Eur J Clin Pharmacol.* 2006; 62(12): 1065-73. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17024485>
- [4] Wilson M-MG, Kaiser FE, Morley JE. Tablet-Breaking Ability of Older Persons With Type 2 Diabetes Mellitus. *Diabetes Educ.* 2001; 27(4): 530-40. Available from: <http://journals.sagepub.com/doi/10.1177/014572170102700408>

## P187

### Adesão ao tratamento em pacientes portadores de esquizofrenia

Almeida BC<sup>1</sup>, Ferreira GE<sup>2</sup>, Souza JN<sup>3</sup>, Gomes LB<sup>4</sup>, Almeida GH<sup>5</sup>

<sup>1</sup>Ciências da saúde, Unichristus, Brasil, beatriz@live.com

<sup>2</sup>Ciências da saúde, Unichristus, Brasil, guilherme\_185@hotmail.com

<sup>3</sup>Ciências da saúde, Unichristus, Brasil, jessicarpb@hotmail.com

<sup>4</sup>Ciências da saúde, Unichristus, Brasil, liviabessagomes@gmail.com

<sup>5</sup>Saúde mental, Universidade Federal do Ceará, Brasil, gilsonholanda@gmail.com

**INTRODUÇÃO:** A esquizofrenia é um distúrbio psicótico grave de ocorrência em todas as populações e culturas e, quando aparece, traz consigo, não raro, efeitos devastadores. Citam-se entre os sintomas mais característicos os transtornos na forma e no conteúdo do pensamento, as alucinações e o afastamento social. A prevalência é cerca de 1% durante toda a vida e acomete preferencialmente adultos jovens, na idade educacional ou início da vida profissional. **OBJETIVO:** Conhecer a percepção dos pacientes portadores de esquizofrenia a respeito de fatores que influenciam na adesão ao tratamento proposto. **MÉTODOS:** Estudo exploratório, transversal, quali-quantitativo. Foram escolhidos de modo aleatório 30 pacientes portadores de transtorno esquizofrênico internados há pelo menos 15 dias em hospital psiquiátrico de referência onde a pesquisa se realizou. Foi utilizado um questionário para conhecer o perfil socioeconômico e um outro composto por quatro questões discursivas, cujas respostas foram gravadas e transcritas, sendo os dados transferidos para o software do Discurso do Sujeito Coletivo (DSC). Houve pré-teste dos instrumentos. O projeto foi aprovado pelo Comitê de Ética em Pesquisas do Centro Universitário Christus (UNICHRISTUS), sob o número 49562915.3.0000.5049. **RESULTADOS:** A amostra foi composta por 63% de homens, 75% solteiros, 53% tinham escolaridade fundamental incompleta, 59% eram católicos, média de idade de 41,6 anos, 13,8 anos em média de doença, 7 recaídas e 5,9 internações ao longo do tempo, renda familiar de 1,3 salários mínimos. A maioria reconhecia que a internação se dera por alteração grave no estado de saúde mental, acreditava estar seguindo o tratamento na internação de modo adequado embora desejasse que os medicamentos fossem diminuídos. Quanto às causas para possível interrupção do tratamento no futuro são apontados a presença de sintomas colaterais, uma possível cura, a falta de suporte familiar ou de eficácia do tratamento e o longo tempo do tratamento. Na maioria das falas, contudo, registra que o tratamento não será interrompido no futuro sob qualquer condição. **CONCLUSÃO:** Os discursos coletados na pesquisa

revelam o conhecimento dos fatores que motivam a não adesão do portador de esquizofrenia ao tratamento, o que se constitui matéria importante a ser incorporada às estratégias terapêuticas e que podem auxiliar na prevenção das recaídas e reinternações.

**Keywords:** *esquizofrenia, adesão, tratamento, discurso do sujeito coletivo, DSC*

#### References:

- [1] Assis JC, Villares CC, Bressan, RA. Entre a razão e a ilusão-desmistificando a loucura. 1th ed. São Paulo: Segmento Farma Editores; 2008.
- [2] Elbogen EB, et al. Medication nonadherence and substance abuse in psychotic disorders: impact of depressive symptoms and social stability. *J Nerv Ment Dis.* 2005; 193(10): 673-79.
- [3] Lefevre F, Lefevre AMC. O discurso do sujeito coletivo: um novo enfoque em pesquisa qualitativa. Caxias do Sul: Educus; 2003.
- [4] Dixon L. Providing services to families of persons with schizophrenia: present and future. *J Mental Health Policy Econ.* 1999; 2(1): 3-8.
- [5] Falkai P, et al. Diretrizes da Federação Mundial das Sociedades de Psiquiatria Biológica para o Tratamento Biológico da Esquizofrenia. Parte I: tratamento.

## P189

### Leisure: formative and well-being space in young people of Viseu

Paulo Eira<sup>1</sup>, António Azevedo<sup>2</sup>, Abel Figueiredo<sup>3</sup>

<sup>1</sup>CI&DEI, Instituto Politécnico de Viseu, Portugal, peira@esev.ipv.pt

<sup>2</sup>CI&DEI, Instituto Politécnico de Viseu, Portugal, toazevedo@esev.ipv

<sup>3</sup>CI&DEI, Instituto Politécnico de Viseu, Portugal, abel.figueiredo@esev.ipv.pt

**BACKGROUND:** Health and well-being are associated in an ineluctable way. Predictor of well-being and, consequently, quality of life, leisure is an important part of healthy life's equation. In this understanding, the concept of health, as well as its experience, can be controlled predominantly by behavioral changes in leisure contexts<sup>[3,8]</sup>. The current research paradigms are found in the inherent processes in subjective well-being and within the scope of positive psychology<sup>[2,4,7]</sup>, using different terminologies, as well as theoretical and conceptual definitions (happiness, affections, emotions, resilience/coping, satisfaction with life), and are determined by social, cultural, psychological and biochemical factors<sup>[6,5,2]</sup>. **OBJECTIVES:** We focus on young's development, centering our work on the knowledge of the offered possibilities so they can live their leisure time in order to obtain healthy life habits. It follows from this objective the analysis of their choices, inferring that individuals who interact effectively with their involvement in various practices, such as sports, music, studies, have positive feelings, feelings of satisfaction. **METHODS:** This research was qualitative in nature, using semi-structured interviews applied to a study group composed of young people who attend secondary school in the municipality of Viseu. The content analysis was developed in three phases: pre-analysis (analysis of the obtained information); material's exploitation (in-depth analysis of the obtained information); data processing, interference and interpretation (comparison and analysis of the obtained data)<sup>[1,9]</sup>. **RESULTS:** It was inferred that young people show positive levels of well-being when involved in physical and sports practices. The boys, when in competitive sport practices and the girls, when in sports and leisure recreational practices. **Discussion:** Physical activities can be understood as means to satisfy new individual needs, to achieve an active and dynamic lifestyle that enhances values and attitudes. **CONCLUSIONS:** Sport practice is considered an added value that binds for the pleasure associated to its practices,