

OC13: Cytotoxicity of the soil fungus *Pisolithus tinctorius* spores against cancer cells – a potential source of anticancer drugs

Ricardo Alves¹, João Fernandes¹, Marco Preto², Rui S. Oliveira^{1,3,4}, Rosário Martins^{1,2}

¹Research Centre on Health and Environment, School of Allied Health Technologies, Polytechnic Institute of Porto, Vila Nova de Gaia, Portugal

²CIIMAR - Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Porto, Portugal

³Centre for Functional Ecology, Department of Life Sciences, University of Coimbra, Portugal

Presenting author: ricardoalvest@gmail.com

Introduction: Cancer is one of the major causes of death worldwide being the search for new anticancer drugs essential for the treatment of this disease. The sporocarps of the soil fungus *Pisolithus tinctorius* contain pisosterol, a triterpene that has been shown to have antitumor activity against some cancer cell lines. Nevertheless, no studies have focused on the anticancer potential of other structures such as spores, and so the anticancer potential of *P. tinctorius*, remains largely unknown.

Objectives: The main objective of this study was to evaluate the potential of *P. tinctorius* spores as a source of anticancer compounds.

Materials and Methods: A crude extract of spores was prepared with Dichloromethane/Methanol (DCM/MeOH-2:1). From the crude extract, 11 fractions were prepared with increasing polarity. The viability of the colon adenocarcinoma cell line RKO, breast carcinoma cell line T47D, osteocarcinoma cell line MG63 and the normal cell hCMEC/D3 exposed to crude extract and fractions was assessed by the 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide reduction assay.

Results and Discussion: The concentration test of 0.1 mg/ml showed a reduction in cell viability of around 95% in all cancer cell lines in the fractions D, F, G and H, without significant reduction in viability of hCMEC/D3 cells.

Conclusion: *P. tinctorius* spores exert cytotoxic activity in cancer cell lines, while having little effects on normal cells, which highlights their anticancer potential.

Acknowledgements: R.S. Oliveira acknowledges the support of Fundação para a Ciência e a Tecnologia (FCT) through research grant SFRH/BPD/85008/2012 and Fundo Social Europeu. This work was financed by national funds through FCT within the scope of the Project PEst-C/MAR/LA0015/2013.

References

1. Montenegro, R. C., Andrade-Neto, M., Elisabete, M., Pessoa, C. & Costa-Lotufo, L. (2004). Cytotoxic activity of pisosterol, a triterpene isolated from *Pisolithus tinctorius* (Mich.: Pers.) Coker & Couch, 1928. Zeitschrift für Naturforschung C, 59, 519-5.