

Poster 6

In vitro antibacterial and cytotoxic activity of *Laurobasidium lauri* extracts

L. Silva ¹, A. I. Oliveira ², F. Moreira ², R. Ferraz Oliveira ² and C. Pinho ^{2,*}

¹ Escola Superior de Saúde, Instituto Politécnico do Porto, R. Dr. António Bernardino de Almeida 400, 4200-072 Porto, Portugal

² Centro de Investigação em Saúde e Ambiente (CISA), Escola Superior de Saúde, Instituto Politécnico do Porto, R. Dr. António Bernardino de Almeida 400, 4200-072 Porto, Portugal

* Correspondence: clp@ess.ipp.pt

Abstract

Background: Plants and mushrooms have been used as medicines for many years, as a source of antibiotics, antineoplastics, among others [1]. Research have been conducted on the medicinal uses of Portuguese plants, however, the therapeutic potentials of some of these plants used in traditional medicine, has remained unexploited. **Objective:** This study aims to evaluate antibacterial and cytotoxic activities, *in vitro*, of *Laurobasidium lauri*, a well-known fungus used in folk medicine on Madeira Island. **Methods:** An experimental study was performed using two extracts (aqueous and ethanolic 55% (V/V)) of the fungus isolated and in combination with three medicinal plants (*Parietaria judaica*, *Polygonum aviculare*, and *Peperomia galioides*). The antibacterial activity was evaluated against *Escherichia coli* and *Staphylococcus aureus*, through disc diffusion and broth microdilution methods. Cytotoxicity was evaluated using MTT (3-[4,5-dimethylthiazol-2-yl]-2,5 diphenyl tetrazolium bromide) assay. **Results:** Regarding the disc diffusion method *E. coli* was not susceptible to any of the extracts, except for the antibiotic (41.0 ± 1.0 mm). However, *S. aureus*, when subjected to 10 mg/ml of the ethanolic extracts of *L. lauri* (isolated and in combination) exhibited an inhibition with a diameter of 16,3 ± 2,5 mm and 12,0 ± 1,7 mm, respectively, when compared to the control, ciprofloxacin (24,0±1,0 mm). Also, the ethanolic extract of the isolated fungus had the best value of minimum inhibitory concentration (MIC) for *S. aureus* (MIC = 0.078125 mg/mL). The ethanolic extract of the fungus in combination with medicinal plants showed greater cytotoxic action on lung cancer cells A549 (IC₅₀ = 48.3±1.0 µg/mL). **Conclusions:** The fungus presented cytotoxic and antibacterial potential, and the results observed may be related to some bioactive compounds (e.g., costunolide and dehydrocostuslactone, two natural sesquiterpene lactones present in *Laurus* trees, where the fungus grows) [2]. However, more research is needed to confirm these biological activities and mechanisms of action.

Keywords: *Laurobasidium lauri*; antibacterial activity; cytotoxicity

Acknowledgments

This research received no external funding.

References

1. Friedman, M. Mushroom polysaccharides: chemistry and antiobesity, antidiabetes, anticancer, and antibiotic properties in cells, rodents, and humans. *Foods* **2016**, *5*, 80.
2. Ferrari, B.; Castilho, P.; Tomi, F.; Rodrigues, A. I.; do Ceu Costa, M.; Casanova, J. Direct identification and quantitative determination of costunolide and dehydrocostuslactone in the fixed oil of *Laurus novocanariensis* by 13C-NMR spectroscopy. *Phytochem Anal* **2005**, *16*, 104-107.



In *Scientific Letters*, works are published under a CC-BY license (Creative Commons Attribution 4.0 International License at <https://creativecommons.org/licenses/by/4.0/>), the most open license available. The users can share (copy and redis-

tribute the material in any medium or format) and adapt (remix, transform, and build upon the material for any purpose, even commercially), as long as they give appropriate credit, provide a link to the license, and indicate if changes were made (read the full text of the license terms and conditions of use at <https://creativecommons.org/licenses/by/4.0/legalcode>).