

III 1H-TOXRUN
INTERNATIONAL
CONGRESS
2024

No Boundaries for
Toxicology: One Health,
One Society, One Planet

The Big Challenges of
the 21st Century

2-3 May 2024 | Porto, Portugal

SCIENTIFIC LETTERS

BOOK OF ABSTRACTS

Volume 1 Supplement 1 (2024)

ISSN 2795-5117

Poster 35

Antioxidant profile of Portuguese and Spanish craft beers

Maria João Pereira^{1,*}, Diana Santos¹, Cláudia Pinho² and Ana Isabel Oliveira²

¹ ESS, Polytechnic of Porto, Rua Dr. António Bernardino de Almeida, 400 4200 - 072, Porto, Portugal

² LAQV/ REQUIMTE, ESS, Polytechnic of Porto, Rua Dr. António Bernardino de Almeida, 400, 4200-072 Porto, Portugal

* Correspondence: psj@ess.ipp.pt

<https://doi.org/10.48797/sl.2024.246>

Abstract

Background: The antioxidant potential of craft beer (CB) may be due to high quality of the raw materials (water, malt, hop and yeast) and traditional techniques [1-4]. The scarcity of studies evaluating the antioxidant potential of CB highlights the relevance of this study. **Objective:** *In vitro* evaluation of the antioxidant activity of aqueous extracts of Portuguese and Spanish CB. **Methods:** Experimental study using six CB with different styles: Milk Stout (EL-MS), India Pale Ale (EL-IPA, ALM-IPA), Imperial Stout (EME-IS), Oatmeal Stout (ALM-OS), Pilsner (EL-P), Munich Dunkell (B-MD) and two industrial beers, Pilsner (S-P) and Munich Dunkell (S-MD). The pH, acidity content (AC) and total phenolic compounds (TPC) were determined. The antioxidant capacity was assessed using 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) and the hydrogen peroxide (H₂O₂) assays, expressed in concentration for 50% activity inhibition (IC₅₀). One-way ANOVA and Student's t-test were used for statistical analysis in GraphPad[®] Prism 8.0 software, with a significance level of 0.05. **Results:** The pH of the beers varied between 3.89±0.00-4.78±0.05 and AC between 0.13±0.01-0.44±0.01%. ALM-IPA had the highest TPC value (8.96±0.64mg gallic acid equivalents/g). IPA style presented the lower IC₅₀ values in H₂O₂ (ALM-IPA with an IC₅₀=23.54±1.53µg/mL, *p*<0.05) and ABTS (EL-IPA with an IC₅₀=55.21±4.68µg/mL, *p*<0.05) assays. The industrial beers have lower TPC values compared to CB (same style, *p*<0.05), and lower capacity to neutralize the H₂O₂ and ABTS radicals. For Omisore *et al.* (2005), samples with IC₅₀>50µg/mL are classified as being moderately active, while samples with IC₅₀<50µg/mL have high antioxidant capacity. Also, the results for IPA style are in line with Breda *et al.* (2022), in which light-colored beers had better antioxidant profiles. However, according to Silva *et al.* (2022) the best antioxidant profiles were associated with dark beers. **Conclusions:** The samples showed antioxidant potential, but further tests should be carried out considering the complex underlying antioxidant mechanisms.

Keywords: craft beer; antioxidant activity; phenolic compounds

Acknowledgments

This research received no external funding.

References

1. DRE. Portaria n.º 91/2022 de 9 de fevereiro. Diário da República (2022), 1ª Série, 4–7.
2. Villacreses, S.; Blanco, C.A.; Caballero, I. Developments and characteristics of craft beer production processes. *Food Biosci* (2022), 45, 1–17.
3. Silva, S.; Oliveira, A.; Cruz, A.; Oliveira, R.; Almeida, R.; Pinho, C. Physicochemical properties and antioxidant activity Portuguese craft beers and raw materials. *Molecules* (2022), 27, 1–15.
4. Breda, C.; Barros, A.I.; Gouvinhas, I. Characterization of bioactive compounds and antioxidant capacity of Portuguese craft beers. *Int J Gastron Food Sci* (2022), 27, 1–7.
5. Omisore, N.O.A.; Adewunmi, C.O.; Iwalewa, E.O.; Ngadjui, B.T.; Adenowo, T.K.; Abegaz, B.M.; Ojewole, J.A.; Watchueng, J. Antitrichomonal and antioxidant activities of *Dorstenia barteri* and *Dorstenia convexa*. *Brazilian J Med Biol Res* (2005), 38, 1087–1094.